


REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE Nov 2000		3. REPORT TYPE AND DATES COVERED Proceedings of Environmental Ergonomics VIII OCT 1998
4. TITLE AND SUBTITLE WOMEN LITTER CARRIERS: OBSERVATIONS AND MODELING			5. FUNDING NUMBERS	
6. AUTHOR(S) W.R. SANTEE AND W.T. MATTHEW				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE NATICK, MA 01760-5007			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. ARMY MEDICAL RESEARCH & MATERIEL COMMAND FORT DETRICK, MD 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>Combat support roles for female soldiers expose them to extremes of both environment and physical exertion. Our goals were to quantify female performance under extreme conditions and to evaluate using male-based models to predict female performance. We collected data on 4 female soldiers performing tasks related to litter casualty evacuation in a hot-dry environment. Activities consisted of walking (1.34 m×s-1) and self-paced two person litter carriage (68 kg load) to simulate casualty evacuation. Uniforms were worn with (MOPP-4) and without (MOPP-0) Chemical Protective (CP) clothing and masks. There were statistically significant differences (p=0.006) between MOPP-0 and MOPP-4 for rectal temperature (DTre) while walking, but not for litter carriage. The primary limiting factor for litter carriage was muscular-skeletal strain. Without CP clothing, thermal strain was not the immediate limiting factor. Data for Tre were compared to values calculated with the Heat Strain Decision Aid (HSDA) and SCENARIO models. The Root Mean Squared Deviations (RMSD) from the observed Tre values were compared to the mean standard deviations ('S'D'). Both male-based models were applicable to the female subject population. The HSDA demonstrated better fit, but SCENARIO is more suitable for accurate and direct input of intermittent work.</p> <div style="text-align: center;">  <span style="font-size: 2em; font-weight: bold;">20001124 056</span> </div>				
14. SUBJECT TERMS Keywords: modeling, chemical protective clothing, load carriage, women, heat			15. NUMBER OF PAGES 4	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT U	18. SECURITY CLASSIFICATION OF THIS PAGE U	19. SECURITY CLASSIFICATION OF ABSTRACT U	20. LIMITATION OF ABSTRACT U	

# Environmental Ergonomics VIII

International Series on Environmental Ergonomics, Volume 1.

---

Editors: James A. Hodgdon  
Jay H. Heaney  
Michael J. Buono

Naval Health Research Center  
and  
San Diego State University  
San Diego, California, USA



Selected papers from the  
8th International Conference on Environmental Ergonomics

San Diego, California, USA

18-23 October 1998

# WOMEN LITTER CARRIERS: OBSERVATION AND MODELING

W.R. Santee and W.T. Matthew

U.S. Army Research Institute of Environmental Medicine  
Natick, MA USA



## INTRODUCTION

Combat support roles for female soldiers expose them to extremes of both environment and physical exertion. We investigated tasks related to the evacuation of casualties by litter. Our goals were to quantify female performance under extreme conditions and to compare the efficacy of using male-based models to predict female performance.

## MATERIALS AND METHODS

Data were collected on 4 female soldiers in a hot-dry environment (1). Investigators adhered to AR 70-25 and USAMRDC Regulation 70-25 on Use of Volunteers in Research. Subjects participated in these studies after giving their free and informed voluntary consent. Test activities consisted of walking ( $1.34 \text{ m}\cdot\text{s}^{-1}$ ) forward to the casualty and self-paced 2-person litter carriage (68 kg load) to simulate stages in casualty evacuation. Metabolic costs for rest, walking and litter carriage were measured prior to testing. Litter carriage was intermittent work consisting of a series of short carries separated by rest periods. Clothing consisted of the Battledress Uniform (BDU) or the Battledress Overgarment (BDO). The BDU provided no chemical protection (MOPP-0 condition), whereas the BDO was worn with the M-40 mask (MOPP-4 condition). Data for rectal temperature ( $T_{re}$ ) were compared to values calculated with the Heat Strain Decision Aid (HSDA) (2) and SCENARIO (3) models. The HSDA model is an executable version of the USARIEM Heat Strain Model (4) developed to predict soldier performance. Model inputs are air temperature, relative humidity, wind speed, solar radiation, clothing, activity or metabolic rate and values for soldier height and weight and heat acclimation. The SCENARIO model was developed as a post hoc model to evaluate experimental results and requires additional inputs of maximum oxygen uptake and globe or mean radiant temperature. Although SCENARIO allows direct input of time-indexed metabolic rates, the HSDA does not. Consequently, metabolic inputs for the HSDA consisted of weighted average values that included a time adjustment to compensate for the metabolic cost of each lift.

## RESULTS

Table 1 presents mean meteorological data used as modeling inputs, and Table 2 summarizes the test results. Mean values for subject height and weight were respectively  $162 \pm 1 \text{ cm}$  and  $63 \pm 5 \text{ kg}$ . There were statistically significant differences between MOPP-0 and MOPP-4 for  $\Delta T_{re}$  ( $P = 0.006$ ) while walking, but not for litter carriage. The primary limiting factor for litter carriage was muscular-skeletal stress.

**Table 1.** Environmental conditions during walking (W) and litter carriage (L) test activities in MOPP-0 (BDU) or MOPP-4 (BDO) clothing

Test	Meteorological values (SD)						
	$T_a$ °C	$T_g$ °C	wind $m \cdot s^{-1}$	RH %	WBGT °C	$T_{rad}$ °C	$T_{grnd}$ °C
W-BDU	39.41 (0.43)	51.11 (0.96)	2.49 (0.98)	12.57 (1.00)	28.9 (0.5)	82.20 (6.87)	52.95 (1.30)
W-BDO	37.96 (0.27)	47.05 (0.44)	4.82 (1.25)	17.85 (0.52)	28.3 (0.3)	84.05 (4.72)	50.52 (0.21)
L-BDU	37.82 (0.34)	46.42 (0.65)	4.63 (1.13)	17.72 (1.57)	28.1 (0.4)	81.12 (4.86)	49.80 (1.17)
L-BDO	35.46 (0.33)	42.21 (1.98)	4.82 (1.10)	26.79 (1.10)	27.4 (0.8)	71.53 (7.30)	46.38 (1.60)

**Table 2.** Summary of subject responses by activity<sup>1</sup>

Test	$T_{re}$ , °C	$\Delta T_{re}$ , °C	Endurance time in minutes	
			range	mean
W-BDU	38.33 (0.42)	0.93 (0.27)	80-160 <sup>2</sup>	140 (40)
W-BDO	38.55 (0.26)	1.28 (0.40)	20-37	29 (8)
L-BDU	37.86 (0.50)	0.69 (0.51)	28-107	59 (35)
L-BDO	38.00 <sup>3</sup> (0.50)	0.76 <sup>3</sup> (0.41)	22-57 <sup>3</sup>	43 <sup>3</sup> (19)

<sup>1</sup> Values in parentheses are Standard Deviations

<sup>2</sup> Maximum time for any activity was 160 min

<sup>3</sup> Three subjects, otherwise n=4

Figures 1 and 2 plot the outputs from both models with mean  $T_{re}$  values. The Root Mean Squared Deviations (RMSD) (5) from the observed  $T_{re}$  values were compared to the mean standard deviations (SD). For all days and models (4x2), with one exception,  $RMSD \leq SD$ . Overall, the HSDA model fit the data better. For litter carriage, the 2-day average RMSD for HSDA was 0.06 vs. 0.10 for SCENARIO.

## DISCUSSION

With the exception of walking with chemical protection (MOPP-4),  $T_{re}$  values remained below 38.5°C. This indicates that factors other than thermal strain were limiting performance. Figures 1 and 2 show that both models project the general response patterns, although  $T_{re}$  is clearly overpredicted for walking in MOPP-4. When HSDA was used to model male performance during another study (6), the rate of increase in core temperature was overpredicted. It may be

desirable for a predictive model to provide a conservative estimate of core temperature, and thus provide some margin for error. The comparison of modeling results to this limited data set suggests there may be no such margin for error when the model is run with inputs for a female population.




Figure 1. Comparison of observed mean  $T_{re}$  to modeling results for walking in MOPP-0 (A) and MOPP-4 (B)




Figure 2. Comparison of observed mean  $T_{re}$  to modeling results for litter carriage in MOPP-0 (A) and MOPP-4 (B)

## CONCLUSIONS

The results indicate that strength and equipment are limiting factors for litter carriage regardless of chemical protection (CP) clothing or weather condition. Without CP clothing, thermal strain was not the immediate limiting factor. For modeling, an accurate estimate of metabolic cost of each activity is critical. Both male-based models were applicable to our female subject population. The HSDA demonstrated better fit, but the format of SCENARIO is more suitable for direct input of metabolic rates for intermittent work.

## REFERENCES

1. Santee, W.R., Kraning, K.K. and Matthew, W.T. 1997, Modeling analysis of female litter bearers during heat stress, *Aviation, Space and Environmental Medicine*, **68**, 647.
2. Joppa, M.D. 1993, *P2NBC2 Heat Strain Decision Aid: User's Guide Version 2.1.*, (San Diego, CA:Scientific Applications International Corporation).
3. Kraning, K.K. and Gonzalez, R.R. 1997, A mechanistic computer simulation of human work in heat that accounts for physical and physiological effects of clothing, aerobic fitness, and progressive dehydration, *Journal of Thermal Biology*, **22**, 331-342.
4. Pandolf, K.B., Stroschein, L.A., Drolet, L.L., Gonzalez, R.R. and Sawka, M.N. 1986, Prediction modeling of physiological responses and human performance in heat, *Computers in Biology and Medicine*, **16**, 310-329.
5. Haslam R.A. and Parsons K.C. 1989, Models of human response to hot and cold environments, Final Report Vol. 1, Report No: APRE 2170/106, Army Personnel Research Establishment, Farnborough, UK.
6. Santee, W.R., Matthew, W.T. and Tharion, W.J. 1992, Simulated approach marches during thermal stress: A P2NBC2 study, Technical Report T12-92, U.S. Army Research Institute of Environmental Medicine, Natick, MA.

## ACKNOWLEDGEMENT AND DISCLAIMER

The study was funded by the Defense Women's Health Research program. The views expressed in this report are those of the authors and should not be construed as official Department of the Army position, policy or decision, unless so designated by other official documentation.